CASE SHEETS

Name

:Unnikrishnan.

Age

:29 years.

Sex

:Male.

Address

:Mappatuthodi house,

Iravimangalam,

Perunthalmannu

Cause of the burn:

- ✓ Dry heat.
- Moist heat.
- Chemicals.
- Electric burns.
- Radiation burns.

Degree of the burn:

FIRST DEGREE	SECOND DEGREE	THIRD DEGREE
Redness	 Blister formation 	 Painless
Severe pain	✓ Sepsis	 Destruction of skin
 Singering of hair 	Severe pain	 Disfigurement
 Burning sensation 	 Raise in temperature 	 Disablement
 Raise in temperature 	 Discolouration 	

Depth of the burn:

- Epidermal burns.
- ✓ Dermo epidermal burns.
- Deep burns.

Thickness of the burn:

- · Full thickness burns.
- ✓ Partial thickness burns.

Surface area of the burn:

	Area	Percentage
	ricad and neck	9%
1	Chest and abdomen	18%
	Back	18%
1	Right upper arm	9%
1	Left upper arm	9%
	Right lower arm	10/0
	Left lower arm	18%
	Perineum	1%

Total percentage of burns:

10%

Prognosis:

First degree-curable.

Second degree-curable with disfigurement.

Third degree-incurable.

Systemic examination:

Central Nervous System Fear for fire. Cardio Vascular System Normal. Respiratory System Normal.

Urinary System Burning micturition.

Reproductive System Normal.

History of the burn:

The lamp turned and the fire caught in the shirt. In the effort to quench the fire he got injured in the hands and stomach. Was hospitalised for a month. Visited the hospital on alternate days and did dressing. The wound healed after 2 months.

Presenting complaints:

Wound is healed and thick keloid scar tissue formed along with white discolouration in the hands and stomach

Duration :4 months

Treatment history:

6-12-99

Was admitted in Thrissur Jubilee Hospital for a month and came to Pathanjali Ayurveda Hospital after 4 months. The medications are as follows.

11-6-99	Manjishtadi ks+guggulu panchaphala
	[15ml bat 45 ml warm water 6 and]

[15ml ks+45 ml warm water, 6 am]

Chandrapraba pill. [ks+pill,6 pm]

Aragwadarishtam+Kumariasavam.

[30 ml,3:30/8:30 pm]

Agnijith.

[External application]

Thickness of the keloid scar tissue reduced. 25-6-99

16-7-99 White discolouration reduced. Skin has become soft. 3-9-99

Very good improvement.

Manjishtadi ks+guggulu panchaphala choornam.

[15 ml ks+45 ml warm water]

Mandoora vadakam pill+buttermilk.

[1:30 pm] Aragwadarishtam.

[30 ml ks, 8:30 pm]

Normal skin colour and texture attained.

Result of the treatment:

The scarring and discolouration is removed by treatment in a duration of 6 months.

Advice during the course of treatment:

- The wound should not be exposed to any liquids.
- 2. Intake of Non-vegetarian foods should be avoided.
- The wound should not be bandaged.