Name

Salsal

Age

: I year. :Male.

Address

S/o Mr. Mansoor,

Pothanchery, Parayapuram.

Cause of the burn

- Dry heat.
- ✓ Moist heat.
- Chemicals.
- Electric burns.
- Radiation burns.

Degree of the burn:

FIRST DEGREE		SECOND DEGREE		THIRD DEGREE	
-	Redness		Blister formation		Painless
V	Severe pain	V	Sepsis		Destruction of skin
	Singering of hair		Severe pain		Disfigurement
1	Burning sensation		Raise in temperature		Disablement
-	Raise in temperature		Discolouration		

Depth of the burn:

- ✓ Epidermal burns.
- Dermo epidermal burns.
- Deep burns.

Thickness of the burn:

- Full thickness burns.
- ✓ Partial thickness burns.

Surface area of the burn:

Area	ercentage
 Head and neck 	9%
✓ Chest and abdomen	18%
Back	18%
✓ Right upper arm	9%
✓ Left upper arm	9%
 Right lower arm 	18%
 Left lower arm 	18%
Perineum	1%

Total percentage of burns:

8%

Prognosis:

✓ First degree-curable.

Second degree-curable with disfigurement.

Third degree-incurable.

Systemic examination

Central Nervous System Cardio Vascular System

Fear for everything. Increased pulse rate.

Respiratory System Urinary System Reproductive System

Normal. Normal.

History of the burn:

When playing, a vessel containing hot water turned over him and got injured.

Presenting complaints:

White discoloured patches on chest, stomach and hands. Wound not healed in the right ankle joint. Fever and shivering present.

Duration

:1 month.

Treatment history:

Was hospitalised in Kozhikode medical college for 15 days and then continued the treatment at home. Arrived at Pathanjali Ayurveda Hospital after a month. The medicaments were,

24-5-99

Aragwadarishtam+saribadiasavam+

gorochanadi pill. [30 ml,1:30/8:30 pm]

Agnijith.

[External application]

31-5-99 Granulation tissue formed in the right ankle joint.

9-6-99 Healing of the wound is completed.

Swelling, discolouration and itching present.

Aragwadarishtam+Kumariasavam.

[30 ml, 1:30/8:30 pm]

9-7-99 Swelling disappeared, discolouration fading, slight itching present.

Aragwadarishtam+loha sindooram 21

[30 ml, 1:30/8:30 pm]

23-10-99 Itching absent, slight scarring present.

Scar tissue has become soft and the skin regaining its normal

texture.

14-3-00 Skin has become normal.

Result of the treatment:

The second degree burn was completely cured in a duration of 12 months and was advised to apply Agnijith for another 6 months.

Advice during the course of treatment:

- 1. The wound should not be exposed to any liquids.
- Intake of Non-vegetarian foods should be avoided.
- 3. The wound should not be bandaged.